



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
28.04.1999 Bulletin 1999/17

(51) Int Cl.⁶: **C12N 15/54, C12N 9/12,
C12N 1/21, C12N 5/10**

(21) Application number: **98307747.0**

(22) Date of filing: **24.09.1998**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• **Akira, Shizuo**
Minoo-shi, Osaka 562-0031 (JP)
• **Kawai, Taro**
Nishinomiyashi, Hyogo 663-8136 (JP)

(30) Priority: **26.09.1997 JP 261589/97**

(74) Representative: **Kremer, Simon Mark et al**
Mewburn Ellis,
York House,
23 Kingsway
London WC2B 6HP (GB)

(71) Applicant: **Japan Science and Technology
Corporation**
Kawaguchi-shi, Saitama 332-0012 (JP)

(54) **DNA coding for serine/threonine kinase**

(57) There is provided a DNA coding for a serine/
threonine kinase.

Thus, the present invention provides a DNA coding
for a protein (a) or (b):

(a) a protein comprising the amino acid sequence
as shown in SEQ ID NO: 1;

(b) a protein comprising an amino acid sequence
having one or several amino acids deleted, substi-
tuted or added in the amino acid sequence as
shown in SEQ ID NO: 1, and exhibiting a serine/
threonine kinase activity.

BEST AVAILABLE COPY

Description

BACKGROUND OF THE INVENTION

5 Field of the Invention:

[0001] The present invention relates to a serine/threonine kinase, a DNA coding for said kinase, a recombinant vector comprising said DNA, a transformant transformed with said vector, and a process for preparing the serine/threonine kinase.

10

Prior Art:

15

[0002] Various signals from the exterior of a cell are transmitted through receptors on the cell surface into the cell and ultimately into the nucleus. The signals transmitted into the nucleus activate transcription factors and, as a result, expression of a group of genes is induced or repressed to produce phenotypes such as cell proliferation, differentiation and cell death. Many transcription factors have been cloned and the structure of functional domains have been elucidated: MOLECULAR BIOLOGY OF THE CELL THIRD EDITION, pp. 401-469. These functional domains are known to include leucine zipper, helix-loop-helix and zinc finger structures. Among them, the leucine zipper structure is a motif commonly found in such transcription factors as Jun/Fos, ATF/CREB and C/EBP families and these transcription factors form homo- or hetero dimers through their leucine zipper structures to control the transcription of specific genes: Hai, T. et al., Proc. Natl. Acad. Sci., USA, 88:3720-3724 (1991).

20

[0003] Recently, it is reported that the leucine zipper structure is also found other functional molecules than the transcription factors (Holzman, L.B. et al., J. Biol. Chem., 269:30808-30817, 1994), suggesting that the leucine zipper structure not only facilitates the binding between transcription factors but also acts generally as a protein-protein interactional domain in cells.

25

[0004] Therefore, identification of molecules interacting with the leucine zipper domain is considered to be useful in analyzing not only new functions of transcription factors but also functions of the leucine zipper structure in other molecules than the transcription factors.

30

SUMMARY OF THE INVENTION

[0005] It is an object of the present invention to provide a serine/threonine kinase, a DNA coding for said kinase, a recombinant vector comprising said DNA, a transformant transformed with said vector, and a process for preparing the serine/threonine kinase.

35

[0006] As a result of their eager studies based on the above described problems, the present inventors have succeeded in isolating a DNA coding for a serine/threonine kinase from cDNA libraries prepared from human placenta and mouse brain and thus completed the present invention.

[0007] Accordingly, the present invention is the following recombinant protein (a) or (b):

40

- (a) a protein comprising the amino acid sequence as shown in SEQ ID NO: 1;
- (b) a protein comprising an amino acid sequence having one or several amino acids deleted, substituted or added in the amino acid sequence as shown in SEQ ID NO: 1, and exhibiting a serine/threonine kinase activity.

45

[0008] Also, the present invention is the following recombinant protein (c) or (d):

- (c) a protein comprising the amino acid sequence as shown in SEQ ID NO: 2;
- (d) a protein comprising an amino acid sequence having one or several amino acids deleted, substituted or added in the amino acid sequence as shown in SEQ ID NO: 2, and exhibiting a serine/threonine kinase activity.

50

[0009] Further, the present invention is a DNA coding for said protein. The DNA include, for example, those comprising the base sequence as shown in SEQ ID NO: 3 or 4.

[0010] The present invention is also a recombinant vector comprising said DNA.

[0011] Still further, the present invention is a transformant transformed with the recombinant vector.

55

[0012] Finally, the present invention is a process for producing a serine/threonine kinase comprising cultivating the transformant in a culture medium and collecting the serine/threonine kinase from the resulting culture.

BRIEF DESCRIPTION OF THE DRAWINGS

[0013] The present invention will hereinbelow be described in detail with reference to the drawings attached in which:

- Fig. 1 shows the results of homology search in amino acid sequence;
- Fig. 2 shows the results of homology search in amino acid sequence;
- Fig. 3 is an electrophoretic photograph showing the results of western blot;
- Fig. 4 is a photograph (the form of an organism) showing the results of colony formation in a selective medium;
- Fig. 5 is an electrophoretic photograph showing the results of northern blot;
- Fig. 6 is a photograph (the form of an organism) showing the results of colony formation in a selective medium;
- Fig. 7 is a photograph of NIH3T3 (the form of an organism) showing the form of apoptosis;
- Fig. 8 shows the fraction of LacZ expression cells showing the form of apoptosis;
- Fig. 9 is an electrophoretic photograph showing the kinase activity of ZIP-kinase; and
- Fig. 10 is a photograph (the form of an organism) showing the intracellular localization of ZIP-kinase.

DESCRIPTION OF THE INVENTION

[0014] The recombinant protein according to the present invention, hereinafter also referred to as "ZIP-kinase", is a protein molecule binding to the leucine zipper domain of a transcription factor called ATF4, and has a serine/threonine kinase activity. The ZIP-kinase is a novel nuclear serine/threonine kinase having the leucine zipper structure and has an activity to induce apoptosis. ATF4 is a leucine zipper type transcription factor which binds to cAMP response element (CRE) and belongs to the ATF/CREB family.

[0015] On the other hand, the DNA according to the present invention is obtained from cDNA libraries prepared from human placenta and mouse brain by screening them using so-called yeast two-hybrid system, and codes for ZIP-kinase. Hereinafter, the DNA will also be referred to as "ZIP-kinase DNA".

[0016] The DNA according to the present invention may be cloned in the following manner:

1. Cloning of ZIP-kinase DNA

(1) Preparation of cDNA libraries from human placenta and mouse brain

[0017] Sources of mRNA may include tissues such as human placenta and mouse brain. Established cell lines from these tissues may also be used as the source.

[0018] The mRNA may be prepared by conventional procedures. For instance, total RNA may be obtained by treating the tissue or cell with a guanidine reagent, and poly (A+) RNA (mRNA) may be obtained by an affinity column method using oligo dT-cellulose or poly U-Sepharose on Sepharose 2B as a carrier, or by a batch method. Also, the poly (A+) RNA may further be fractionated by sucrose density-gradient centrifugation.

[0019] The resulting mRNA is used as a template to synthesize a single-stranded cDNA which is in turn used to synthesize a double-stranded cDNA. A recombinant plasmid is prepared from a suitable vector DNA and used to transform Escherichia coli or the like to yield a cDNA library.

[0020] Alternatively, the cDNA library may be commercially available (CLONETECH).

(2) Construction of plasmid pAS2-1

[0021] From the cDNA library obtained in (1) above, a plasmid is prepared for screening for a desired clone.

[0022] Such a plasmid may be obtained by preparing a chimeric DNA by ligating a DNA coding for mouse ATF4 leucine zipper domain (amino acids 298 to 349 in the sequence of ATF4) to a DNA coding for GAL4 DNA binding domain, and ligating the chimeric DNA to bait plasmid pAS2-1.

(3) Screening

[0023] Then said plasmid is used to screen the cDNA library. In the screening, yeast two-hybrid system may be used. The yeast two-hybrid system is an experimental system capable of detecting interaction between proteins in yeast and is capable of screening the library for cDNA of a protein interacting with the desired protein (bait).

[0024] Positive clones may be selected using the growth in a selective medium free of hystidine, tryptophan or leucine and the activity of β -galactosidase.

(4) Determination of base sequence

[0025] The base sequence is determined for the resulting clone. The sequencing may be carried out by any known method such as Maxam-Gilbert method or the dideoxy method and is usually done using an automated base sequencer.

[0026] SEQ ID NOs: 1 and 2 exemplify the amino acid sequence of ZIP-kinase and the base sequence of ZIP-kinase DNA, respectively, according to the present invention. As far as a protein comprising said amino acid sequence has an activity as a serine/threonine kinase, there may be a mutation or variation of deletion, substitution and/or addition of one or several amino acids in said amino acid sequence as shown in SEQ ID NO: 1. For example, a protein having the amino acid sequence as shown in SEQ ID NO: 1 from which the first amino acid methionine has been deleted may also be included in the present invention.

[0027] Herein the serine/threonine kinase activity means an activity of transferring the terminal phosphate group of ATP to a certain amino acid (serine or threonine) of a protein. The introduction of mutation or variation may be carried out by any known method (Deng, W.P. et al., Anal. Biochem., 200: 81, 1992) or using a commercially available kit (Site-Directed Mutagenesis Kit of CLONETECH).

[0028] Once the base sequence of ZIP-kinase DNA according to the present invention is determined, ZIP-kinase DNA according to the present invention may then be obtained by chemical synthesis, or by PCR with various tissues-derived cDNA as a template, or hybridization of a DNA fragment having said base sequence as a probe.

2. Construction of recombinant vector and transformant

(1) Construction of recombinant vector

[0029] The recombinant vector of the present invention may be obtained by ligating or inserting ZIP-kinase DNA of the present invention into an appropriate vector. The vector for inserting ZIP-kinase DNA of the present invention is not particularly limited as long as it can be replicated in a host, and may include plasmid DNA, phage DNA, etc. The plasmid DNA may be prepared from *E. coli* or *Agrobacterium* by the alkali extraction (Bimboim, H.C. & Doly, J., (1979) Nucleic acid Res., 7:1513) or modified method. Further, commercially available plasmids may also be used, for example, pUC18 (Takara Shuzo), pUC19 (Takara Shuzo), pBluescript SK⁺ (Stratagene), pGEM-T (Promega), pT7 Blue (Novagen) and PBR322 (Takara Shuzo).

[0030] The phage DNA may include, for example, M13mp18, M13mp19, λ gt10, λ gt11, etc.

[0031] When the DNA of the present invention is inserted into a vector, the purified DNA may first be cut with a suitable restriction enzyme and inserted into a restriction enzyme site or multi cloning site of a suitable vector DNA to ligate with the vector.

[0032] The DNA of the present invention should be incorporated into a vector such that the function of the DNA can be realized. In addition to a promoter and the DNA of the present invention, the vector of the present invention may comprise a terminator, a ribosome-binding sequence and the like. The terminator may be a stop codon such as TGA, TAG or TAA and the ribosome-binding sequence may be a leader sequence.

(3) Preparation of transformant

[0033] The transformant of the present invention may be obtained by introducing the recombinant expression vector of the present invention into a host such that the desired gene can be expressed therein.

[0034] The host is not particularly limited so long as the DNA of the present invention can be expressed and may include, for example, bacteria belonging to the genus *Escherichia* or *Bacillus*, such as *Escherichia coli* and *Bacillus subtilis*; yeast, such as *Saccharomyces cerevisiae* and *Saccharomyces pombe*; animal cells, such as COS and CHO cells; and insect cells, such as Sf9.

[0035] When a bacterium, such as *E. coli*, is used as a host, it is preferred that the vector of the present invention is capable of autonomously replicating in said bacterium and comprises a promoter, a ribosome-binding sequence, the DNA of the present invention, and a transcription terminating sequence. The vector may also comprise a gene controlling the promoter.

[0036] For example, pET and pGEX (Pharmacia) may be used as the expression vector.

[0037] Any promoter may be used so long as the expression can be effected in the host such as *E. coli*. A promoter derived from *E. coli* or phage, such as *trp*, *lac*, PL or PR promoter, may be used. An artificially designed and modified promoter, such as T7 or T3 may also be used.

[0038] The method for introducing the recombinant vector into a bacterium is not particularly limited so long as a DNA can be introduced into a bacterium. For example, the method using calcium ion (Proc. Natl. Acad. Sci., USA, 69, 2110-2114 (1972)) and the electroporation method may be used.

[0039] When a yeast is used as a host, YEp13, YEp24 and YCp50 may be used as the expression vector. The

promoter used is not particularly limited so long as the expression in the yeast can be effected, and may include, for example, gall, ga110, heat shock protein, MF α 1 and SV40 promoters.

[0040] The method for introducing the recombinant vector into a yeast is not particularly limited so long as a DNA can be introduced into a yeast, and include, for example, the electroporation method (Methods Enzymol., 194, 182-187 (1990)), the spheroplast method (Proc. Natl. Acad. Sci., USA, 84, 1929-1933 (1978)), and the lithium acetate method (J. Bacteriol., 153, 163-168 (1983)).

[0041] When an animal cell is used as a host, an expression vector, such as pcDNA1/Amp or pcDNA1 (Invitrogen) is used. The promoter used may also be the early gene promoter of human cytomegalovirus.

[0042] The method for introducing the recombinant vector into an animal cell may include, for example, the electroporation method, the calcium phosphate method and the lipofection method.

[0043] The recombinant vectors of the present invention (one vector containing ZIP-kinase DNA from human placenta and another vector containing ZIP-kinase DNA from mouse brain) have been introduced into E. coli DH5, E. coli (hZIP-kinase) DH5 and E. coli (mZIP-kinase) DH5, respectively, and deposited at National Institute of Bioscience and Human-Technology (NIBH), Agency of Industrial Science and Technology, 1-3, Higashi 1-chome, Tsukuba-shi, Ibaraki-ken 305, Japan on September 25, 1997 under Accession Nos. FERM BP-6487 and FERM BP-6488, respectively.

3. Production of ZIP-kinase

[0044] ZIP-kinase of the present invention may be obtained by cultivating the transformant in a culture medium and collecting from the resulting culture.

[0045] The transformant of the present invention may be cultivated in a culture medium by any method conventionally used to cultivate a host.

[0046] The culture medium for cultivating a transformant obtained from a microorganism such as E. coli or yeast as a host may be either a natural or synthetic medium so long as it contains a carbon source, a nitrogen source and inorganic salts which can be utilized by the microorganism and the transformant can efficiently be cultivated.

[0047] The carbon source used may include carbohydrates, such as glucose, fructose, sucrose, starch, and dextrose; organic acids, such as acetic acid and propionic acid; and alcohols, such as ethanol and propanol.

[0048] The nitrogen source which may be used includes ammonia; ammonium salts of inorganic or organic acids, such as ammonium chloride, ammonium sulfate, ammonium acetate and ammonium phosphate; other nitrogen containing compounds; peptone, meat extract, corn steep liquor, and yeast extract.

[0049] The minerals which may be used include monobasic potassium phosphate, dibasic potassium phosphate, magnesium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, copper sulfate, calcium carbonate, calcium chloride, and disodium phosphate.

[0050] The cultivation is generally carried out at 37°C for 12 to 18 hours under aerobic conditions, such as shaking culture and aerated spinner culture. During the cultivation, pH is kept at 7.0 to 7.5. The pH is adjusted with an inorganic or organic acid or alkaline solution, or carbonic acid gas.

[0051] During the cultivation, an antibiotic such as ampicillin or tetracycline may optionally be added to the medium.

[0052] When a microorganism transformed with an expression vector comprising an inducible promoter is cultivated, an inducer may be added to the medium, if necessary. For example, when a microorganism transformed with an expression vector comprising Lac promoter is cultivated, isopropyl- β -D-thiogalactopyranoside (IPTG) or the like may be added to the medium. When a microorganism transformed with an expression vector comprising trp promoter is cultivated, indole-acrylic acid (IAA) may be added to the medium.

[0053] When a transformant obtained from an animal cell as a host is cultivated, a conventional culture medium such as RPMI 1640 or DMEM medium or these media to which fetal bovine serum is added may be used.

[0054] The cultivation is generally carried out at 37°C for 1 to 3 days in the presence of 5% CO₂.

[0055] During the cultivation, an antibiotic such as kanamycin or penicillin may be added to the medium.

[0056] After the cultivation, when ZIP-kinase of the present invention is produced in the host cell, the ZIP-kinase is extracted by disruption of the cell. When ZIP-kinase of the present invention is produced in the exterior of the cell, the culture may be directly used as it is, or the ZIP-kinase of the present invention may be isolated and purified from the culture, after removing the cell by centrifugation, using any conventional biochemical methods generally used in the isolation and purification of proteins, such as ammonium sulfate precipitation, gel chromatography, ion exchange chromatography and affinity chromatography, singly or in any combination thereof.

EXAMPLES

[0057] The present invention will be further illustrated by the following examples. However, the scope of the present invention is not limited to these examples.

Example 1: Cloning of ZIP-kinase DNA

(1) Preparation of cDNA library

5 [0058] A commercially available cDNA (CLONETECH) was used in the present invention.

(2) Construction of plasmid

[0059] DNA coding for leucine zipper domain of mouse ATF4 was obtained by PCR method.

10 [0060] The composition of the PCR reaction was 1.0 µg DNA, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl₂, 0.2 mM dNTP, 0.5 µM primer, and 1 U Taq.

[0061] The following primers were used:

15 Sense primer: 5' -GGGAATTCGCGGAGCAGGAGGCT - 3' (SEQ ID NO: 5)

Antisense primer: 5' -GGGGATCCCTAGGGGACCCTTTTCTA - 3' (SEQ ID NO:

20 6)

[0062] Sense primer: 5'-GGGAATTCGCGGAGCAGGAGGCT-3' (SEQ ID NO: 5) Antisense primer: 5'-GGGGATCCCTAGGGGACCCTTTTCTA-3' (SEQ ID NO: 6)

25 [0063] PCR reaction was first carried out at 94°C for 1 minute. Then, 25 cycles of reactions at 94°C for 20 seconds, at 56°C for 20 seconds and at 72°C for 30 seconds were carried out. Finally, the reaction at 72°C for 10 minutes was effected.

30 [0064] The PCR products were cut with EcoRI/BamHI and inserted into EcoRI/BamHI site of pAS2-1 vector. The plasmids were used to transform E. coli DH5 α and purified by means of a commercially available kit (Wizard miniprep: Promega) based on alkali-SDS method. This plasmid capable of expressing a fused protein of GAL4 DNA binding domain in yeast was used as a bait.

(3) Screening

35 [0065] Yeast strain Y190 was transformed with the plasmid as a bait using MATCHMAKER Two-Hybrid System kit of CLONETECH. Transformants were selected by growth in tryptophan(-) medium as an index. Further, cDNA libraries capable of expressing a fusion protein with GAL4 transcription activating domain (CLONETECH, mouse brain and human placenta MATCHMAKER cDNA libraries) were transformed. Transformants can grow in tryptophan(-), leucine (-) medium. Further, since reporter genes, HIS3 and LacZ genes, were transcribed if the bait bound to the DNA coding for the protein from the library, positive clones can grow in tryptophan (-), leucine (-), histidine (-) medium and provides blue color in the presence of X-gal because of their β -galactosidase activity. Plasmids were purified from the positive clones using MATCHMAKER Two-Hybrid System kit of CLONETECH, and used to transform E. coli. Plasmids were purified from the resulting transformants and the base sequences thereof were determined (ABI model 377). The resulting base sequences were searched for homology using GenBank, EMBL, DDBJ data base.

40 [0066] As a result, those having high homology (20% or higher) with the previously reported C/EBP family, AP-1 family and genes having leucine zipper structure were identified as novel genes. Seven (7) and 2 clones of such genes were obtained from mouse brain and human placenta cDNA libraries, respectively. All these genes were derived from an identical gene.

(4) Determination of base sequence

50 [0067] The thus obtained gene was considered to code for a kinase and the DNA coding for this novel kinase was designated as ZIP-kinase DNA (Zipper Interacting Protein Kinase DNA). The base sequence of the full length ZIP-kinase DNA was determined.

55 [0068] The base sequences of ZIP-kinase DNA obtained from human placenta and mouse brain are shown in SEQ ID NOs: 3 and 4, respectively. The amino acid sequences encoded by the base sequences of SEQ ID NOs: 3 and 4 are shown in SEQ ID NOs: 1 and 2, respectively.

[0069] The amino acid sequence encoded by ZIP-kinase DNA obtained from human placenta (human ZIP-kinase) and the amino acid sequence encoded by ZIP-kinase DNA obtained from mouse brain (mouse ZIP-kinase) were

searched for homology therebetween and the leucine zipper domain and serine/threonine kinase domain were found in the C- and N-terminal of the respective amino acid sequences, respectively (Fig. 1). Further, mouse and human ZIP-kinases consisted of 448 and 454 amino acids, respectively, and the homology between mouse and human was 84.9% at amino acid level.

[0070] Moreover, the kinase domains of the ZIP-kinases showed high homology with DAP-kinases positively controlling apoptosis caused by IFN- γ , suggesting that these kinases form a new family (Fig. 2).

[0071] Example 2: Construction of recombinant vector and preparation of transformant

[0072] To construct a recombinant vector of ZIP-kinase DNA, cDNA coding for ZIP-kinase was synthesized by PCR method.

[0073] The PCR reaction mixture and primers used were as follows.

[0074] The composition of the PCR reaction was 1.0 μ g DNA, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM $MgCl_2$, 0.2 mM dNTP, 0.5 μ M primer, and 1 U Taq.

Sense primer: 5'-GGGTCGACCAC CATGGCTTAC CCATACGATG TTCCAGATTA
CGCTATGTCC ACATTCAGGC AA-3' (SEQ ID NO: 7)

Antisense primer: 5'-GGGTCGACTA GCGCACGCCG CACTCAGCCT GC-3'
(SEQ ID NO: 8)

[0075] PCR reaction was first carried out at 96°C for 1 minute. Then, 30 cycles of reactions at 96°C for 30 seconds, at 56°C for 30 seconds and at 72°C for 1 minute were carried out. Finally, the reaction at 72°C for 10 minutes was effected.

[0076] The resulting PCR products were cut with Sall, inserted into expression vector pEF-BOS (Takara, Ligation kit), and used to transform E. coli DH5 (TOYOBO). Plasmids were purified from the E. coli (Promega, Wizard miniprep) and confirmed by DNA sequence (ABI, model 377).

[0077] A DNA coding for a variant of ZIP-kinase in which the 42nd amino acid lysine in the amino acid sequence as shown in SEQ ID NO: 2 was changed to alanine, hereinafter referred to as "ZIP-kinase K42A", was constructed by using Site-Directed Mutagenesis Kit of CLONETECH. Also, a DNA coding for another variant in which the 422nd and 429th amino acids valine and the 436th amino acid leucine in the amino acid sequence as shown in SEQ ID NO: 2 were changed to alanines, hereinafter referred to as "ZIP-kinase LA", was similarly constructed.

Example 3: Function of DNA of the present invention

(1) Binding of ZIP-kinase to ATF4 in cells

[0078] Whether ZIP-kinase binds to ATF4 in cells as well or not was investigated.

[0079] First, DNA coding for mouse ZIP-kinase (309-448 amino acids in the amino acid sequence as shown in SEQ ID NO: 2) was inserted into expression vector pEF-BOS. Thus, a tag of the transcription factor Myc was provided at the N-terminal end of ZIP-kinase, whereby a DNA coding for Myc-ZIP-kinase complex was designated to have the tag as an epitope to construct the vector (pEF-BOS-Myc-ZIP-kinase). Also, an expression vector (pEF-BOS-FLAG-ATF4) was constructed comprising a DNA coding for human ATF4 (full length)-FLAG complex in which FLAG epitope had been added to the N-terminal end of human ATF4.

[0080] These vectors were transiently introduced into COS-7 cell line by the lipofection method and expressed (Fig. 3 in which lanes 1, 4, 7 and 10 represent FLAG-ATF4; lanes 2, 5, 8 and 11 Myc-ZIP-kinase; lanes 3, 6, 9 and 12 FLAG-ATF4 and Myc-ZIP-kinase). 36 hours after the introduction, the cells were collected and solubilized with 0.5% Nonidet P-40 lysis buffer. The resulting solubilized cell (WCE: whole cell extract) was developed in SDS-PAGE, and transferred to nitrocellulose membrane. Western blot analysis was done using anti-FLAG monoclonal antibody (Fig. 3, lanes 1, 2 and 3) and anti-Myc monoclonal antibody (lanes 7, 8 and 9) to confirm the expression of Myc-ZIP-kinase and FLAG-ATF4.

[0081] Subsequently, the WCE was immunoprecipitated with anti-Myc monoclonal antibody and the precipitate was subjected to the western blot analysis using anti-FLAG monoclonal antibody, attempting to detect co-immunoprecipitation of Myc-ZIP-kinase and FLAG-ATF4 (Fig. 3, lanes 4 to 6).

[0082] As a result, a band of FLAG-ATF4 was detected in lane 6 (Fig 3). For further confirmation, the WCE was then immunoprecipitated with anti-FLAG monoclonal antibody and the precipitate was subjected to the western blot analysis

using anti-Myc monoclonal antibody (lanes 10, 11 and 12). A band of Myc-ZIP-kinase immunoprecipitated with FLAG-ATF4 was found only in lane 12.

[0083] Thus, it was shown that ZIP-kinase and ATF4 bind to each other in cells as well.

[0084] From this result that ZIP-kinase and ATF4 binds to each other, it may be considered that ATF4 may possibly control the activity of ZIP-kinase.

(2) Determination of domain necessary for binding of ZIP-kinase to ATF4

[0085] The site to which ZIP-kinase and ATF4 bind was determined using yeast two-hybrid system. First, variants of mouse ZIP-kinase were prepared: 1) amino acids 278 to 448 of ZIP-kinase (ZIP-kinase 278-448); 2) leucine zipper domain of ZIP-kinase (amino acids 398 to 448) (ZIP-kinase LZ); and 3) a variant of ZIP-kinase in which valine and leucine in the leucine zipper domain were substituted with alanine (ZIP-kinase LA). Each of these variants was designed to produce a chimeric protein with GAL4 trans activating domain, and DNA coding for said chimeric protein was inserted into pACT2 and introduced into yeast strain Y190 together with pAS2-1-ATF4 LZ. The strain was cultivated on histidine+, tryptophan-, leucine-, and histidine-, tryptophan-, leucine- selective media.

[0086] Yeast containing DNA coding for ZIP-kinase 278-448 and yeast containing DNA coding for ZIP-kinase LZ could form a colony on the histidine-, tryptophan-, leucine- medium, indicating that ZIP-kinase bound to ATF4 through leucine zipper domain present at the C-terminal (Fig. 4). Further, when valine and leucine in the leucine zipper domain structure were substituted with alanine, the binding to ATF4 was no longer found.

[0087] Accordingly, it has been elucidated that ZIP-kinase and ATF4 bind to each other through their respective leucine zipper domain.

(3) Expression of ZIP-kinase in each tissue

[0088] Northern blot analysis was carried out to investigate the expression of ZIP-kinase in each tissue.

[0089] As shown in Fig. 5, mRNA of ZIP-kinase (about 1.4 kb) was distributed almost all tissues investigated. However, only low expression was observed in the spleen.

(4) Confirmation of formation of homodimer of ZIP-kinase

[0090] The leucine zipper domain present at the C-terminal of ZIP-kinase is considered to be a domain through which proteins bind to each other. Whether ZIP-kinase forms a homodimer or not was investigated. Plasmid pAS2-1 into which DNA coding for leucine zipper domain of ZIP-kinase was inserted, and plasmid pACT2 into which DNA coding for the leucine zipper domain of ZIP-kinase and a variant in which valine and leucine in said domain were substituted with alanine, were co-introduced into yeast and colony formation was observed in a selective medium.

[0091] As shown in Fig. 6, only yeast co-expressing the ZIP-kinase leucine zipper domain could grow in histidine-, tryptophan-, leucine- medium. Thus, it has been elucidated that ZIP-kinase forms a homodimer through its leucine zipper structure.

(5) Induction of apoptosis by ZIP-kinase

[0092] It has been shown that DAP-kinase, which has high homology with kinase domain of ZIP-kinase, induces apoptosis in HeLa cell. Whether ZIP-kinase has an apoptosis activity was investigated.

[0093] RNA wild type ZIP-kinase tagged with HA (pEF-BOS-HA-ZIP-kinase), a variant thereof in which lysine (42nd amino acid), which is present in ZIP-kinase subdomain II and conserved in other kinases, was substituted with alanine (pEF-BOS-HA-ZIP-kinase K42A), and a variant in which valine and leucine in the leucine zipper domain were substituted with alanine (pEF-BOS-HA-ZIP-kinase LA) were prepared, and DNA coding for each of these proteins was transiently introduced into NIH 3T3 cell together with LacZ expression vector (pEF-BOS-LacZ). After 36 hours from the introduction, X-gal staining was effected.

[0094] As a result, a form of cells stained blue was observed under a microscope (Fig. 7). As compared with a control pEF-BOS-mock (Fig. 7, left, upper), the cell into which the wild type ZIP-kinase was introduced (Fig. 7, right, upper) exhibited a typical form of apoptosis associated with agglomeration of nucleus. The fraction of LacZ expression cells showing the apoptosis form was measured to be 44.9% (Fig. 8).

[0095] On the other hand, such change of form was not observed in the ZIP-kinase-K42A variant (Fig. 7, left, lower) and there was no significant difference in the fraction of apoptosis between the variant and control. Further, in the ZIP-kinase-LA (Fig. 7, right, lower), some cells caused apoptosis but the fraction thereof was significantly reduced as compared with the wild type

[0096] From the above results, the kinase activity of ZIP-kinase is considered to be essential for the induction of

apoptosis by the expression of ZIP-kinase. Further, since apoptosis was suppressed in variants in which a homodimer between ZIP-kinases was inhibited, it is suggested that ZIP-kinases form a homodimer to become an activated form.

(6) Kinase activity of ZIP-kinase

[0097] Whether ZIP-kinase indeed has an activity as a kinase was investigated.

[0098] Each of pEF-BOS-HA-ZIP-kinase, pEF-BOS-HA-ZIP-kinase K42A, and pEF-BOS-HA-ZIP-kinase LA was transiently introduced into COS-7 cell, and 36 hours later, the cell was collected and solubilized with 0.5% Nonidet P-40 lysis buffer. The solubilized cell was immunoprecipitated with anti-HA monoclonal antibody and the kinase activity in the precipitate was detected by in vitro kinase assay (Fig. 9A).

[0099] As a result, a band of phosphorylation by ZIP-kinase was observed at about 50 kDa in the wild type ZIP-kinase (Fig. 9A, lane of HA-ZIP-kinase), while no band corresponding thereto was observed in ZIP-kinase K42A. On the other hand, a phosphorylation band was observed in ZIP-kinase LA. However, when the expression of HA-ZIP-kinase and its variant in the solubilized cell was checked by the western blotting using anti-HA monoclonal antibody, the amount of HA-ZIP-kinase expressed was markedly reduced as compared with the other two variants (Fig. 9B). From this result, it may be considered that the kinase activity observed in ZIP-kinase LA would be very weak as compared with the wild type.

[0100] Further, the amount of wild type ZIP-kinase expressed was low in COS-7 cells; this is considered to be resulted from some lethal effect, such as apoptosis, of ZIP-kinase on COS-7 cells.

(7) Localization of ZIP-kinase in cells

[0101] Knowledge of intracellular localization of ZIP-kinase would be considered to be very effective in analyzing the functions of ZIP-kinase. The present inventors have investigated the localization of ZIP-kinase using a confocal laser microscope.

[0102] An expression vector (pEF-BOS-FLAG-ATF4) comprising DNA coding for ATF4 tagged with FLAG or another vector (pEF-BOS-FLAG-ZIP-kinase K42A) coding for ZIP-kinase K42A tagged with FLAG was transiently introduced into COS-7 cells. After 36 hours, the cells were fixed, reacted with anti-FLAG monoclonal antibody, and stained using FITC-labelled anti-mouse immunoglobulin antibody as a secondary antibody.

[0103] When observed under the confocal laser microscope, the cytoplasm was not stained and the nucleus was stained in the FLAG-ATF4 introduced cells (Fig. 10A). When the localization of FLAG-ZIP-kinase was similarly investigated, the same staining pattern as in ATF4 was observed, confirming that it was localized in the nucleus (Fig. 10B).

[0104] Accordingly, it could be concluded that the ZIP-kinase is a novel nuclear serine/threonine kinase.

Advantages of the Invention:

[0105] According to the present invention, there are provided a serine/threonine kinase, a DNA coding for said kinase, a recombinant vector comprising said DNA, and a transformant transformed with said vector, and a process for the preparation of the serine/threonine kinase.

[0106] Since the ZIP-kinase has a function of inducing apoptosis, the ZIP-kinase and DNA coding for said kinase are useful in being utilizable as a gene therapeutical agent against a cancer and as an anti-cancer agent.

Annex to the description

[0107]

5

SEQUENCE LISTING

10 <110> Japan Science and Technology Corporation

15 <120> DNA CODING FOR SERINE/THREONINE KINASE

<130> PH-569

20

<140>

<141>

25

<150> JP97/261589

<151> 1997-09-26

30

<160> 8

35 <170> PatentIn Ver. 2.0

40 <210> 1

<211> 454

<212> PRT

45

<213> Homo sapiens

<400> 1

50

Met Ser Thr Phe Arg Gln Glu Asp Val Glu Asp His Tyr Glu Met Gly

1

5

10

15

Glu Glu Leu Gly Ser Gly Gln Phe Ala Ile Val Arg Lys Cys Arg Gln

55

EP 0 911 408 A2

	20	25	30
	Lys Gly Thr Gly Lys Glu Tyr Ala Ala Lys Phe Ile Lys Lys Arg Arg		
5	35	40	45
	Leu Ser Ser Ser Arg Arg Gly Val Ser Arg Glu Glu Ile Glu Arg Glu		
10	50	55	60
	Val Asn Ile Leu Arg Glu Ile Arg His Pro Asn Ile Ile Thr Leu His		
	65	70	75
15	80		
	Asp Ile Phe Glu Asn Lys Thr Asp Val Val Leu Ile Leu Glu Leu Val		
	85	90	95
20	100	105	110
	Ser Gly Gly Glu Leu Phe Asp Phe Leu Ala Glu Lys Glu Ser Leu Thr		
	115	120	125
25	130	135	140
	Glu Asp Glu Ala Thr Gln Phe Leu Lys Gln Ile Leu Asp Gly Val His		
	145	150	155
30	160		
	Tyr Leu His Ser Lys Arg Ile Ala His Phe Asp Leu Lys Pro Glu Asn		
	165	170	175
35	180	185	190
	Ile Met Leu Leu Asp Lys Asn Val Pro Asn Pro Arg Ile Lys Leu Ile		
	195	200	205
40	210	215	220
	Asp Phe Gly Ile Ala His Lys Ile Glu Ala Gly Asn Glu Phe Lys Asn		
	225	230	235
45	240		
	Ile Phe Gly Thr Pro Glu Phe Val Ala Pro Glu Ile Val Asn Tyr Glu		
	245	250	255
50			
	Pro Leu Gly Leu Glu Ala Asp Met Trp Ser Ile Gly Val Ile Thr Tyr		
	255		
55			
	Ile Leu Leu Ser Gly Ala Ser Pro Phe Leu Gly Glu Thr Lys Gln Glu		
	260	265	270
60	275	280	285
	Thr Leu Thr Asn Ile Ser Ala Val Asn Tyr Asp Phe Asp Glu Glu Tyr		
	290	295	300
65	305		
	Phe Ser Asn Thr Ser Glu Leu Ala Lys Asp Phe Ile Arg Arg Leu Leu		

EP 0 911 408 A2

[illegible]

450

<210> 2

<211> 448

<212> PRT

<213> Mus musculus

<400> 2

Met Ser Thr Phe Arg Gln Glu Asp Val Glu Asp His Tyr Glu Met Gly

1 5 10 15

Glu Glu Leu Gly Ser Gly Gln Phe Ala Ile Val Arg Lys Cys Gln Gln

20 25 30

Lys Gly Thr Gly Met Glu Tyr Ala Ala Lys Phe Ile Lys Lys Arg Arg

35 40 45

Leu Pro Ser Ser Arg Arg Gly Val Ser Arg Glu Glu Ile Glu Arg Glu

50 55 60

Val Ser Ile Leu Arg Glu Ile Arg His Pro Asn Ile Ile Thr Leu His

65 70 75 80

Asp Val Phe Glu Asn Lys Thr Asp Val Val Leu Ile Leu Glu Leu Val

85 90 95

Ser Gly Gly Glu Leu Phe Asp Phe Leu Ala Glu Lys Glu Ser Leu Thr

100 105 110

Glu Asp Glu Ala Thr Gln Phe Leu Lys Gln Ile Leu Asp Gly Val His

115 120 125

Tyr Leu His Ser Lys Arg Ile Ala His Phe Asp Leu Lys Pro Glu Asn

130 135 140

Ile Met Leu Leu Asp Lys His Ala Ala Ser Pro Arg Ile Lys Leu Ile

145 150 155 160

Asp Phe Gly Ile Ala His Arg Ile Glu Ala Gly Ser Glu Phe Lys Asn

165 170 175

Ile Phe Gly Thr Pro Glu Phe Val Ala Pro Glu Ile Val Asn Tyr Glu

180 185 190

EP 0 911 408 A2

Pro Leu Gly Leu Glu Ala Asp Met Trp Ser Ile Gly Val Ile Thr Tyr
5 195 200 205
Ile Leu Leu Ser Gly Ala Ser Pro Phe Leu Gly Glu Thr Lys Gln Glu
 210 215 220
10 Thr Leu Thr Asn Ile Ser Ala Val Asn Tyr Asp Phe Asp Glu Glu Tyr
225 230 235 240
Phe Ser Ser Thr Ser Glu Leu Ala Lys Asp Phe Ile Arg Arg Leu Leu
15 245 250 255
Val Lys Asp Pro Lys Arg Arg Met Thr Ile Ala Gln Ser Leu Glu His
 260 265 270
20 Ser Trp Ile Lys Val Arg Arg Arg Glu Asp Gly Ala Arg Lys Pro Glu
 275 280 285
25 Arg Arg Arg Leu Arg Ala Ala Arg Leu Arg Glu Tyr Ser Leu Lys Ser
290 295 300
His Ser Ser Met Pro Arg Asn Thr Ser Tyr Ala Ser Phe Glu Arg Phe
30 305 310 315 320
Ser Arg Val Leu Glu Asp Val Ala Ala Ala Glu Gln Gly Leu Arg Glu
 325 330 335
35 Leu Gln Arg Gly Arg Arg Gln Cys Arg Glu Arg Val Cys Ala Leu Arg
 340 345 350
40 Ala Ala Ala Glu Gln Arg Glu Ala Arg Cys Arg Asp Gly Ser Ala Gly
 355 360 365
Leu Gly Arg Asp Leu Arg Arg Leu Arg Thr Glu Leu Gly Arg Thr Glu
45 370 375 380
Ala Leu Arg Thr Arg Ala Gln Glu Glu Ala Arg Ala Ala Leu Leu Gly
385 390 395 400
50 Ala Gly Gly Leu Lys Arg Arg Leu Cys Arg Leu Glu Asn Arg Tyr Asp
 405 410 415
55 Ala Leu Ala Ala Gln Val Ala Ala Glu Val Gln Phe Val Arg Asp Leu

EP 0 911 408 A2

	420	425	430	
5	Val Arg Ala Leu Glu Gln Glu Arg Leu Gln Ala Glu Cys Gly Val Arg			
	435	440	445	
10				
	<210> 3			
	<211> 2132			
15	<212> DNA			
	<213> Homo sapiens			
20				
	<220>			
	<221> CDS			
25	<222> (94)..(1455)			
	<400> 3			
30	gttgccatta ggggactcct gaggtcctat ctccaggctg cggtgactgc actttccctg 60			
	gagtggaagc tgctggaagg cggaccggcc gcc atg tcc acg ttc agg cag gag 114			
35	Met Ser Thr Phe Arg Gln Glu			
	1 5			
	gac gtg gag gac cat tat gag atg ggg gag gag ctg ggc agc ggc cag 162			
40	Asp Val Glu Asp His Tyr Glu Met Gly Glu Glu Leu Gly Ser Gly Gln			
	10	15	20	
	ttt gcg atc gtg cgg aag tgc cgg cag aag ggc acg ggc aag gag tac 210			
45	Phe Ala Ile Val Arg Lys Cys Arg Gln Lys Gly Thr Gly Lys Glu Tyr			
	25	30	35	
50	gca gcc aag ttc atc aag aag cgc cgc ctg tca tcc agc cgg cgt ggg 258			
	Ala Ala Lys Phe Ile Lys Lys Arg Arg Leu Ser Ser Ser Arg Arg Gly			
	40	45	50	55
55	gtg agc cgg gag gag atc gag cgg gag gtg aac atc ctg cgg gag atc 306			

EP 0 911 408 A2

	Val Ser Arg Glu Glu Ile Glu Arg Glu Val Asn Ile Leu Arg Glu Ile	
5	60 65 70	
	cgg cac ccc aac atc atc acc ctg cac gac atc ttc gag aac aag acg	354
	Arg His Pro Asn Ile Ile Thr Leu His Asp Ile Phe Glu Asn Lys Thr	
10	75 80 85	
	gac gtg gtc ctc atc ctg gag ctg gtc tct ggc ggg gag ctc ttt gac	402
	Asp Val Val Leu Ile Leu Glu Leu Val Ser Gly Gly Glu Leu Phe Asp	
15	90 95 100	
	ttc ctg gcg gag aaa gag tcg ctg acg gag gac gag gcc acc cag ttc	450
20		
	Phe Leu Ala Glu Lys Glu Ser Leu Thr Glu Asp Glu Ala Thr Gln Phe	
	105 110 115	
	ctc aag cag atc ctg gac ggc gtt cac tac ctg cac tct aag cgc atc	498
25		
	Leu Lys Gln Ile Leu Asp Gly Val His Tyr Leu His Ser Lys Arg Ile	
	120 125 130 135	
	gca cac ttt gac ctg aag ccg gaa aac atc atg ctg ctg gac aag aac	546
30		
	Ala His Phe Asp Leu Lys Pro Glu Asn Ile Met Leu Leu Asp Lys Asn	
	140 145 150	
35		
	gtg ccc aac cca cga atc aag ctc atc gac ttc ggc atc gcg cac aag	594
	Val Pro Asn Pro Arg Ile Lys Leu Ile Asp Phe Gly Ile Ala His Lys	
	155 160 165	
40		
	atc gag gcg ggg aac gag ttc aag aac atc ttc ggc acc ccg gag ttt	642
	Ile Glu Ala Gly Asn Glu Phe Lys Asn Ile Phe Gly Thr Pro Glu Phe	
	170 175 180	
45		
	gtg gcc cca gag att gtg aac tat gag ccg ctg ggc ctg gag gcg gac	690
	Val Ala Pro Glu Ile Val Asn Tyr Glu Pro Leu Gly Leu Glu Ala Asp	
	185 190 195	
50		
	atg tgg agc atc ggt gtc atc acc tat atc ctc ctg agc ggt gca tcc	738
	Met Trp Ser Ile Gly Val Ile Thr Tyr Ile Leu Leu Ser Gly Ala Ser	
55	200 205 210 215	

EP 0 911 408 A2

	ccg ttc ctg ggc gag acc aag cag gag acg ctc acc aac atc tca gcc	786
5	Pro Phe Leu Gly Glu Thr Lys Gln Glu Thr Leu Thr Asn Ile Ser Ala	
	220 225 230	
	gtg aac tac gac ttc gac gag gag tac ttc agc aac acc agc gag ctg	834
10	Val Asn Tyr Asp Phe Asp Glu Glu Tyr Phe Ser Asn Thr Ser Glu Leu	
	235 240 245	
	gcc aag gac ttc att cgc cgg ctg ctc gtc aaa gat ccc aag cgg aga	882
15	Ala Lys Asp Phe Ile Arg Arg Leu Leu Val Lys Asp Pro Lys Arg Arg	
	250 255 260	
	atg acc att gcc cag agc ctg gaa cat tcc tgg att aag gcg atc cgg	930
20	Met Thr Ile Ala Gln Ser Leu Glu His Ser Trp Ile Lys Ala Ile Arg	
	265 270 275	
	cgg cgg aac gtg cgt ggt gag gac agc ggc cgc aag ccc gag cgg cgg	978
25	Arg Arg Asn Val Arg Gly Glu Asp Ser Gly Arg Lys Pro Glu Arg Arg	
	280 285 290 295	
30	cgc ctg aag acc acg cgt ctg aag gag tac acc atc aag tcg cac tcc	1026
	Arg Leu Lys Thr Thr Arg Leu Lys Glu Tyr Thr Ile Lys Ser His Ser	
	300 305 310	
35	agc ttg ccg ccc aac aac agc tac gcc gac ttc gag cgc ttc tcc aag	1074
	Ser Leu Pro Pro Asn Asn Ser Tyr Ala Asp Phe Glu Arg Phe Ser Lys	
40	315 320 325	
	gtg ctg gag gag gcg gcg gcc gcc gag gag ggc ctg cgc gag ctg cag	1122
	Val Leu Glu Glu Ala Ala Ala Ala Glu Glu Gly Leu Arg Glu Leu Gln	
45	330 335 340	
	cgc agc cgg cgg ctc tgc cac gag gac gtg gag gcg ctg gcc gcc atc	1170
50	Arg Ser Arg Arg Leu Cys His Glu Asp Val Glu Ala Leu Ala Ala Ile	
	345 350 355	
	tac gag gag aag gag gcc tgg tac cgc gag gag agc gac agc ctg ggc	1218
55	Tyr Glu Glu Lys Glu Ala Trp Tyr Arg Glu Glu Ser Asp Ser Leu Gly	

EP 0 911 408 A2

360 365 370 375
5 cag gac ctg cgg agg cta cgg cag gag ctg ctc aag acc gag gcg ctc 1266
Gln Asp Leu Arg Arg Leu Arg Gln Glu Leu Leu Lys Thr Glu Ala Leu
380 385 390
10 aag cgg cag gcg cag gag gag gcc aag ggc gcg ctg ctg ggg acc agc 1314
Lys Arg Gln Ala Gln Glu Glu Ala Lys Gly Ala Leu Leu Gly Thr Ser
395 400 405
15 ggc ctc aag cgc cgc ttc agc cgc ctg gag aac cgc tac gag gcg ctg 1362
Gly Leu Lys Arg Arg Phe Ser Arg Leu Glu Asn Arg Tyr Glu Ala Leu
410 415 420
20 gcc aag caa gta gcc tcc gag atg cgc ttc gtg cag gac ctc gtg cgc 1410
Ala Lys Gln Val Ala Ser Glu Met Arg Phe Val Gln Asp Leu Val Arg
425 430 435
25 gcc ctg gag cag gag aag ctg cag ggc gtg gag tgc ggg ctg cgc 1455
Ala Leu Glu Gln Glu Lys Leu Gln Gly Val Glu Cys Gly Leu Arg
440 445 450
30 taggcgcagt ggggtgggcc agggcccagg acagccggag ctgcggcctgc ggtgggggag 1515
35 ctctctgttg acgtgcgcc tccatcgcc cgggtgcctg tccttgccca gcgccaccag 1575
gctggaggcg gaggggagg agctggagcc agggccgtaa gtgcgcaggc aggggtgggt 1635
gtgggacggg gctgcttctc tacacagcct ctacgtggc ctccaccttc acccctgcat 1695
40 cgtcggtagc cctgggaccc tccaggcagc gtggcctgtg gcaccgtgag ggttgggacc 1755
caccgaggcg cagaggcggc ccgaatgcag ccctggttca ggcccggagg agggtttgcg 1815
ggtagttagc cggacaattc ggcgggggtg tgcctgttgc tgccattagc ccaggaggag 1875
45 gtcgtgggac ggggaggggt ggatggacgg cggacaggca gtccccacgc tgctgggttg 1935
cgccgggctt ggtgggggtct tccactgtgt gcccttctcg ccgaggccgg tccccgggt 1995
50 gtgggggtgcc ctgctgcgga ctctccgcg agcccatcg tcgcgcctgt ggacgcctag 2055
gcaagagcgg ccctctgcag ccaagagaaa taaaatactg gcttccagat aaaaaaaaaa 2115
aaaaaaaaa aaaaaaa 2132
55

5 <210> 4
 <211> 1429
 <212> DNA
 10 <213> Mus musculus

 <220>
 15 <221> CDS
 <222> (10)..(1353)

 20 <400> 4
 ccagccgcc atg tcc aca ttc agg caa gag gat gtt gag gac cat tat gag 51
 25 Met Ser Thr Phe Arg Gln Glu Asp Val Glu Asp His Tyr Glu
 1 5 10
 atg gga gag gag ctt ggc agt ggc caa ttt gcc atc gtg cgc aag tgc 99
 30 Met Gly Glu Glu Leu Gly Ser Gly Gln Phe Ala Ile Val Arg Lys Cys
 15 20 25 30
 35 cag cag aag ggc acg ggc atg gag tat gca gcc aag ttc atc aag aag 147
 Gln Gln Lys Gly Thr Gly Met Glu Tyr Ala Ala Lys Phe Ile Lys Lys
 35 40 45
 40 cgg cgc ctg cca tcc agc cgg cgc ggt gtg agc cgg gag gag atc gaa 195
 Arg Arg Leu Pro Ser Ser Arg Arg Gly Val Ser Arg Glu Glu Ile Glu
 50 55 60
 45 cgc gag gtg agc atc ctg cgc gag atc cgc cac ccc aac atc ata aca 243
 Arg Glu Val Ser Ile Leu Arg Glu Ile Arg His Pro Asn Ile Ile Thr
 65 70 75
 50 ctg cat gac gtg ttc gag aac aag aca gat gtg gtg ctg atc ctg gag 291
 Leu His Asp Val Phe Glu Asn Lys Thr Asp Val Val Leu Ile Leu Glu
 55 80 85 90

EP 0 911 408 A2

	ctg gtg tcc ggt ggc gag ctt ttc gac ttc ctg gcc gag aag gag tca	339
5	Leu Val Ser Gly Gly Glu Leu Phe Asp Phe Leu Ala Glu Lys Glu Ser	
	95 100 105 110	
	ttg acg gag gat gag gcc acg cag ttc ctc aaa caa atc cta gac ggt	387
10	Leu Thr Glu Asp Glu Ala Thr Gln Phe Leu Lys Gln Ile Leu Asp Gly	
	115 120 125	
	gtc cac tac ctg cac tcc aag cgc atc gca cac ttt gac ctg aag ccc	435
15	Val His Tyr Leu His Ser Lys Arg Ile Ala His Phe Asp Leu Lys Pro	
	130 135 140	
	gag aac atc atg ttg ctg gac aag cac gca gcc agc ccc cgc att aag	483
20	Glu Asn Ile Met Leu Leu Asp Lys His Ala Ala Ser Pro Arg Ile Lys	
	145 150 155	
	ctc atc gac ttt ggc atc gcg cac agg atc gag gct ggc agc gag ttc	531
25	Leu Ile Asp Phe Gly Ile Ala His Arg Ile Glu Ala Gly Ser Glu Phe	
	160 165 170	
	aag aac atc ttt ggc aca ccc gag ttt gtc gcc ccc gag atc gtg aac	579
30	Lys Asn Ile Phe Gly Thr Pro Glu Phe Val Ala Pro Glu Ile Val Asn	
	175 180 185 190	
	tat gag cca ctt ggc ttg gag gct gac atg tgg agc att ggc gtc atc	627
	Tyr Glu Pro Leu Gly Leu Glu Ala Asp Met Trp Ser Ile Gly Val Ile	
40	195 200 205	
	acc tac atc ctc ctg agc gga gcg tcc cca ttc ctg ggc gag acc aag	675
	Thr Tyr Ile Leu Leu Ser Gly Ala Ser Pro Phe Leu Gly Glu Thr Lys	
45	210 215 220	
	cag gag acg ctg acg aac atc tca gca gtg aac tat gac ttt gat gag	723
50	Gln Glu Thr Leu Thr Asn Ile Ser Ala Val Asn Tyr Asp Phe Asp Glu	
	225 230 235	
	gaa tac ttc agc agc acc agc gag ctg gcc aag gac ttc atc cgc agg	771
55	Glu Tyr Phe Ser Ser Thr Ser Glu Leu Ala Lys Asp Phe Ile Arg Arg	

EP 0 911 408 A2

	240	245	250	
5	ctg ctg gtc aaa gac ccc aag agg agg atg acc atc gca cag agc ctg			819
	Leu Leu Val Lys Asp Pro Lys Arg Arg Met Thr Ile Ala Gln Ser Leu			
	255	260	265	270
10	gag cat tcc tgg atc aag gtg cgc agg cgc gag gac ggc gcc cgg aag			867
	Glu His Ser Trp Ile Lys Val Arg Arg Arg Glu Asp Gly Ala Arg Lys			
	275	280	285	
15	cca gag cga cgg cgg ctg cgc gcc gcg cgc ctg cgc gag tac agc ctc			915
	Pro Glu Arg Arg Arg Leu Arg Ala Ala Arg Leu Arg Glu Tyr Ser Leu			
	290	295	300	
20	aag tcc cac tcg agc atg ccg cgc aac acg agc tac gcc agc ttc gag			963
	Lys Ser His Ser Ser Met Pro Arg Asn Thr Ser Tyr Ala Ser Phe Glu			
	305	310	315	
25	cgc ttc tca cgc gtg ctg gag gac gtg gcg gcg gca gag cag ggg ctg			1011
	Arg Phe Ser Arg Val Leu Glu Asp Val Ala Ala Ala Glu Gln Gly Leu			
30	320	325	330	
	cgc gag ctg cag cga ggc agg cgc cag tgc cgg gag cgc gtg tgt gcg			1059
35	Arg Glu Leu Gln Arg Gly Arg Arg Gln Cys Arg Glu Arg Val Cys Ala			
	335	340	345	350
	ctg cgc gcg gcc gcc gag cag cgg gag gcg cgc tgc cgc gac ggg agc			1107
40	Leu Arg Ala Ala Ala Glu Gln Arg Glu Ala Arg Cys Arg Asp Gly Ser			
	355	360	365	
45	gca ggg cta ggg cgc gac ctg cga cgc ctg cgc acg gag ctg ggg cgc			1155
	Ala Gly Leu Gly Arg Asp Leu Arg Arg Leu Arg Thr Glu Leu Gly Arg			
	370	375	380	
50	acc gag gct ctg cgc acg cgc gcg cag gag gag gcg cgg gcg gcg ctg			1203
	Thr Glu Ala Leu Arg Thr Arg Ala Gln Glu Glu Ala Arg Ala Ala Leu			
	385	390	395	
55	ttg ggt gcc ggg ggc ctg aag cgt cgc ctg tgt cgc ctg gag aac cgt			1251

EP 0 911 408 A2

Leu Gly Ala Gly Gly Leu Lys Arg Arg Leu Cys Arg Leu Glu Asn Arg
 5 400 405 410
 tac gac gcg cta gcc gct cag gtg gcc gct gag gtg caa ttc gtg cgc 1299
 Tyr Asp Ala Leu Ala Ala Gln Val Ala Ala Glu Val Gln Phe Val Arg
 10 415 420 425 430
 gac ctg gtg cgt gcg ctg gag cag gaa cgg ctg cag gct gag tgc ggc 1347
 Asp Leu Val Arg Ala Leu Glu Gln Glu Arg Leu Gln Ala Glu Cys Gly
 15 435 440 445
 gtg cgc taggctgcgg caccgccaga ccccgaccga ccccagaat aaagctgctt 1403
 Val Arg
 20 tccacgtaaa aaaaaaaaaa aaaaaa 1429
 25
 <210> 5
 <211> 23
 30 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 <223> Description of Artificial Sequence:Synthetic
 40 oligonucleotides
 45
 <400> 5
 gggaattcgc ggagcaggag gct 23
 50
 <210> 6
 <211> 26
 55 <212> DNA

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence:Synthetic

10

oligonucleotides

<400> 6

15

ggggatccct aggggaccct ttctta

26

20

<210> 7

<211> 63

25

<212> DNA

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence:Synthetic

35

oligonucleotides

<400> 7

40

gggtcgacca ccatggctta cccatacgat gttccagatt acgctatgtc cacattcagg 60

caa

63

45

<210> 8

<211> 32

50

<212> DNA

<213> Artificial Sequence

55

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotides

<400> 8

gggtcgacta gcgcacgccg cactcagcct gc

32

Claims

1. A recombinant protein (a) or (b):

- (a) a protein comprising the amino acid sequence as shown in SEQ ID NO: 1;
- (b) a protein comprising an amino acid sequence having one or several amino acids deleted, substituted or added in the amino acid sequence as shown in SEQ ID NO: 1, and exhibiting a serine/threonine kinase activity.

2. A recombinant protein (c) or (d):

- (c) a protein comprising the amino acid sequence as shown in SEQ ID NO: 2;
- (d) a protein comprising an amino acid sequence having one or several amino acids deleted, substituted or added in the amino acid sequence as shown in SEQ ID NO: 2, and exhibiting a serine/threonine kinase activity.

3. A DNA coding for a protein (a) or (b):

- (a) a protein comprising the amino acid sequence as shown in SEQ ID NO: 1;
- (b) a protein comprising an amino acid sequence having one or several amino acids deleted, substituted or added in the amino acid sequence as shown in SEQ ID NO: 1, and exhibiting a serine/threonine kinase activity.

4. A DNA coding for a protein (c) or (d):

- (c) a protein comprising the amino acid sequence as shown in SEQ ID NO: 2;
- (d) a protein comprising an amino acid sequence having one or several amino acids deleted, substituted or added in the amino acid sequence as shown in SEQ ID NO: 2, and exhibiting a serine/threonine kinase activity.

5. The DNA of claim 3 comprising the base sequence as shown in SEQ ID NO: 3.

6. The DNA of claim 4 comprising the base sequence as shown in SEQ ID NO: 4.

7. A recombinant vector comprising a DNA of any one of claims 3 to 6.

8. A transformant transformed with a recombinant vector of claim 7.

9. A process for producing a serine/threonine kinase comprising cultivating a transformant of claim 8 in a culture medium and collecting the serine/threonine kinase from the resulting culture.

FIG. 1

mouse ZIP-kinase	MSTFRQEDVEDHYEMGEELGSGQFATVRKCKQKGTGMEYAAKFKKRRLLPSSRRGVSREE 	60
human ZIP-kinase	MSTFRQEDVEDHYEMGEELGSGQFATVRKCRQKGTGKEYAAKFKKRRLLSSRRGVSREE 	60
mouse ZIP-kinase	IEREVSTLREIRHPNIITLHDVFENKTDVLLILELVSGGELFDFLAEKESL TEDEATQFL 	120
human ZIP-kinase	IEREVNITLREIRHPNIITLHDIFENKTDVLLILELVSGGELFDFLAEKESL TEDEATQFL 	120
mouse ZIP-kinase	KQILDGVHYLHKKRIAHFDLKPENIMLLDKHAASPRIKLIDFGIAHRIEAGSEFKNIFGT 	180
human ZIP-kinase	KQILDGVHYLHKKRIAHFDLKPENIMLLDKNVPNPRIKLIDFGIAHRIEAGSEFKNIFGT 	180
mouse ZIP-kinase	PEFVAPEIVNYEPLGLEADWMSIGVITYILLGASPFGETKQETLTNISAVNYDFDEEY 	240
human ZIP-kinase	PEFVAPEIVNYEPLGLEADWMSIGVITYILLGASPFGETKQETLTNISAVNYDFDEEY 	240
mouse ZIP-kinase	FSSTSELAKDFIRRLLVKDPKRRMTIAQSLHSHWIK-VRRR-----EDGARKPPERRLRAA 	295
human ZIP-kinase	FSNTSELAKDFIRRLLVKDPKRRMTIAQSLHSHWIKAIRRRNVRGEDSGRKPERRRLKTT 	300
mouse ZIP-kinase	RLREYSLKSHSSMPRNTSYASFERFSRVLEDVAAAEQGLRELQRGRQCRERVCAALRAAA 	355
human ZIP-kinase	RLKEYTIKSHSLPPNNSYADFERFSKVL EEAEEGLRELQSRRLCHEDVEALAAIY 	360
mouse ZIP-kinase	EQREARCRDGSAGLGRDLRRLRTELGRTEALRTRAQEEAALLGAGGLKRRLCRLNRY 	415
human ZIP-kinase	EEKEAWYREESDSLGDQLRRLRQELLKTEALKRQAEAKGALLGTSGLKRFRSRLNRY 	420
mouse ZIP-kinase	DALAAQVAAAEVQFVRDLVRALEQERLQA-ECGVR 	448
human ZIP-kinase	EALAKQVASEMRVQDLVRALEQEKLGQVEGCLR 	454

Protein kinase domain

Protein kinase domain

Leucine zipper domain

26

FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9A




FIG. 9B




FIG. 10A

FLAG-ATF4





FIG. 10B

FLAG-ZIP-kinase K42A



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.